

Chlorine-Containing Sesquiterpenes of Higher Plants

VALERY M. DEMBITSKY¹ and GENRICH A. TOLSTIKOV²

¹School of Pharmacy,
P. O. Box 12065, The Hebrew University of Jerusalem, Jerusalem 91120 (Israel)

E-mail: dvalery@cc.huji.ac.il

²N. N. Vorozhtsov Novosibirsk Institute of Organic Chemistry, Siberian Branch of the Russian Academy of Sciences, Pr. Lavrentyeva 9, Novosibirsk 630090 (Russia)

E-mail: gtolstik@nioch.nsc.ru

(Received January 21, 2002)

Abstract

The structures of chlorine-containing sesquiterpenes produced by higher plants and data on the biological activity of some compounds are presented.

Contents

Introduction	363
Sesquiterpenic lactones	363
Indanonic sesquiterpenes	367
Other types of sesquiterpenes	369

INTRODUCTION

Sesquiterpenes as a class of terpenoids are widespread in nature. They are produced by plants, microorganisms and inhabitants of the world ocean [1]. Sesquiterpenes containing haloid atoms have begun to be studied comparatively recently. The data available nowadays witness to the fact that halogenized sesquiterpenes, unlike halomonoterpenes and haloditerpenes produced almost exclusively by marine organisms [2, 3], are found among the metabolites of higher plants. In the present review, the structures and partly the biological activity of chlorine-containing sesquiterpenes isolated from plants widespread in nature [4] are considered.

SESQUITERPENIC LACTONES

The first chlorine-containing sesquiterpenoids were eupachlorin (**1**), its acetate (**2**) and eupachloroxin (**3**) isolated in 1968 from the plant *Eupatorium rotundifolium* [5, 6]. All the three compounds have supplemented the series of sesquiterpenic lactones characterized by a high anticancerogenic activity.

The genus *Centaurea* belongs to perennial grasses widespread in Eurasia, North America and Africa. More than 500 species of this genus are known. In West Siberia, *Centaurea adpressa* Ledeb., *C. Cyanus* L., *C. Jaeea* L., *C. Scabiosa* L. and *C. Sibirica* L. grow.

Plants of the genus *Centaurea* have turned out to be producers of a whole group of chlo-

rine-containing sesquiterpenic lactones. As one can see from the structures presented herein below, sesquiterpenic lactones from plants of the genus *Centaurea* belong to Guianic structural type. Their characteristic property is that they belong to esters.

In this connection, the lactones of *Centaurea* may be conditionally divided into three groups depending on the position of the chlorine atom. So, lactones of group 1 contain the chlorine atom in the guanic skeleton. In lactones of group 2, the chlorine atom is situated in the acid substituent. Sesquiterpenes of group 3 are esters of chlorine-containing guanolides and chlorine-containing oxyacids.

Centaurea repens which is poisonous to cattle contains centaurepensin (**4**) whereby its toxicity is mainly accounted for [7, 8]. The same lactonoester has been isolated from the plant *C. solstitialis* [9]. The mentioned plant *C. repens* produces also chlororepdiolide (**5**) [10].


Lactone acroptilin (**6**) or chlorhyssopifolin C has been found in plants *C. repens*, *C. hyrcanica* [11, 12] and *C. hyssopifolium* [13–15].

Extracts from the plants *C. hyssopifolium* [14] and *C. linifolia* [16] contain chlorhyssopifolins B (**7**), D (**8**) and E (**9**). The plant *C. linifolia* is also a producer of linichlorins A (**10**), B (**11**) and C (**12**) [16].


Salegin (**13**) isolated from the plant *Saussereea elegans* [17] is an epimer of linichlorin C. Solstitiolide (**14**) and episolstitiolide (**15**) have been isolated from the plant *C. solstitialis* [9].

C. janeri and *C. nigra* growing in Latin America synthesize the already mentioned lactone (**4**) and chlorjanerin (**16**) [18].


C. aegyptica growing in North Africa contains guanolides (**4**), (**7**), (**9**), (**10**), (**14**) and (**15**) [19]. The Argentinian species of *C. repens*, apart from the above mentioned lactones (**5**), (**10**) and (**16**), produces also trihydroxylactone repensolide (**17**) [20].




12




13




14. R = CO



15. R = CO





In South America there is a plant *C. scoparia* which is a source of lactones (**7**), (**10**), (**16**) and of cebellin D (**18**), its acetate (**19**) and a lactone of abnormal structure (**20**) which carbon skeleton may be formally considered as belonging to *tris*-homosesquiterpenoid or *bis*-norditerpenoid types [21–23].

The species *C. sinaica* widespread in North Africa and Middle East produces lactones (**4**) and (**5**) [24, 25]. In the same regions, the species *C. bella* grows which contains, besides the already mentioned lactones (**4–6**), (**16**) and (**19**), lactone cebellin J (**21**). The latter compound (**21**), just like lactones (**10**), (**14**) and (**15**), is synthesized by the plant *C. adriatica* growing on the shores of the Adriatic Sea [26–28].

In other species of genus *Centaurea*, the above mentioned lactones have also been found. So, *C. hermanii* contains compounds (**10**) and (**16**) [29], *C. imperialis* produces lactones (**4**) and (**19**) [30], from *C. incana* three lactones – (**6**), (**14**) and (**15**) [31] – have been isolated, and *C. marshaliana* contains compounds (**6**), (**16**) and (**18**) [26].

Plants *C. kotschy*, *C. phaeopappoides* and *C. thracica* are interesting for containing the only lactone (**11**) [32].

Plants of genera *Chartolepsis*, *Leuzea* and *Psephellus* growing in Eurasia are related to plants of genus *Centaurea*. This is confirmed by identity of their metabolites to lactones produced by plants of this genus.

So, plants *Chartolepsis bieberstenii*, *Ch. Glasifolia* and *Ch. Pterocaula* produce as the main components lactones (**4**), (**6**) and (**16**), and as minor ones, lactones (**12**) and (**21**) [28].

Three species – *Leuzea carthamoides*, *L. rhabontica* ssp. *helenifolia* and *L. rhabonti-*

coides – contain chlorjanerin (**16**) and an abnormal lactone (**20**) [28].

Eight species of genus *Psephellus* – *P. carthanicus*, *P. colchicus*, *P. dagestanicus*, *P. deae-batus*, *P. karabaghensis*, *P. nogmovii*, *P. somcheticus* and *P. zangezuri* – contain lactones (**4**), (**6**) and (**11**), and three species – *P. declinatus*, *P. hypoleucus* and *P. leucophyllus* – are sources of lactone linichlorin B (**11**) [28, 33].


Perennial plants of genus *Saussurea* growing both in Europe and in Asia contain chlorinated lactones. For example, *Saussurea candicans* produces the above mentioned lactones (**4**), (**10**) and (**16**) [34]. The latter of the three mentioned lactones has been isolated from the Mongolian species *S. liphshitzii* [35], whereas *S. elegans* contains compounds (**10**) and (**13**) [17].

The plant *Ambrosia maritima* is used in the traditional medicine of Egypt [36]. From the leaves of this plant, 11- β -hydroxy-13-chloro-11,13-dihydrochymenine (**22**) has been isolated which displays a high cytotoxic activity with respect to cancerous cells P-388, L-1210, KB and V-79 [37]. Noteworthy is the fact that the lactone (**22**) differs from other lactones in the configuration of angular carbon atoms.

A study of two related plant species *Cousinia canescens* and *C. piptocephala* has demonstrated that they synthesize the known lactones (**10**) and (**16**), and lactone (**25**) [38]. In extracts of leaves of *Gutenbergia marginata* (Vernoniceae), angelate (**23**) and dihydrochloro-angelate (**24**) have been detected [39].

From the Middle Asis plant *Achillea biebersteinii*, biesanine (**26**) has been isolated [40].

Brazilian population of the plant *Lasiolaena santonii* contains ketolactones (**27–29**) char-



acterized by the presence of a dienone group [41]. In Brazil, the plant *Lasiolaena morii* producing, besides the lactone (**28**), also lactone (**30**) which molecule contains such a specific fragment as an hydroxy group at C 15, has been found [42].

It is noteworthy that the only chlorine-containing sesquiterpene contained in the leaves of Brazilian plant *Trichogonia gerdneri* is acetate (**31**) [43].


The same structural peculiarity is characteristic of lactones (**32**) and (**33**) isolated from



the leaves of Costa Rican plant *Mikania viti-folia* [44].

Among six sesquiterpenic lactones produced by the Mexican wormwood species *Artemisia klotzchiana*, one – chloroklotzchin (**34**) – contains chlorine [45].



Epimer epoxylactones – dihydroxybievani-nes (**35**) and (**36**) – have been isolated from the medicinal plant *Chrysanthemum parthenium* [46].

The medicinal plant *Ambrosia maritima* widespread in Egypt synthesizes chloroketolactone chlorambrosine (**37**) [47].




32. R = COC(CH₃)OHCH₂Cl33. R = COC(CH₃)=CH₂


34

35. R₁ = OH, R₂ = CH₃36. R₁ = CH₃, R₂ = OH

37



38



39

An interesting component of the Argentinian species of the plant *Stevia sanguinea* is the cross-conjugate ketolactone (38) [48].

Leaves of plant *Kaunia lasiophthalma* contain 19 sesquiterpenic lactones one of which (39) contains a chlorine atom [49].

The 13-chlorosolstictiphilin (40) and its acetate (41) isolated from the Portuguese plant *Cynara humilis* differ from all the earlier described guianolides by the presence of a chlorine atom at C₁₃ [50].


Chlorovenidol (42) and chlorovenidin (43) are produced by the grassy plant *Vinidium fastuosum* widespread in Egypt [51]. These compounds differ from the above mentioned guianolides by the position of the lactone fragment.

Isolation of chlorine-containing hermacranolides has been described. Epoxy lactone

poskephnolide (44) is produced by the plant *Critoniopsis huaircajana* growing in Costa Rica [52]. Diepoxyde (45) has been identified in extracts from the North American plant *Liatris acidota* [53]. Derivates of hermacrane containing an oxide fragment C₁₁–C₁₄ form a new type of sesquiterpenic lactones referred to heliangolides. Chlorine-containing heliangolides (46) and (47) have been isolated from the plant *Calea morri* [54], whereas the plant *Calea villosa* produces lactones (48) and (49) [55].


INDANONIC SESQUITERPENES

Indanonic sesquiterpenes (50–58) were discovered more than 30 years ago [56]. They have been found in Chinese and Japanese ferns *Pte-*




40. R = H

41. R = Ac



42. R = H

43. R = COCH₂CH₃



Pteridium aquilinum var. *latiusculum* which are toxic to cattle. The indanonic sesquiterpenes themselves display a cytotoxic effect and inhibit the development of sea urchin embryos and infusoria [57].

Pterosine F (**50**) has been detected in many species of fern: *Pteridium* var. *latiusculum* [56–59], *Pteridium* subsp. *wightianum* [60], *Microlepia strigosa* and *M. substrigosa* [61], and in species belonging to the genus *Pteris*: *P. angustifolia*, *P. cretica*, *P. dactylina*, *P. multifida* and *P. tremula* [62].

Pterosine H (or hypolepine A) (**51**) has been found in plants *Dennstaedtia scaba* [63], *Hypolepis punctata* [64], *Pteridium aquilinum* [59],


Pteridium subsp. *wightianum* [60], *Microlepia substrigosa*, *M. obtusiloba*, *M. trapeziformis* and *M. speluncae* [61].


Pterosine J (**52**) has been isolated from two fern species – *Pteridium aquilinum* var. *latiusculum* [58, 65] and *Pteris tremula* [62].

Pterosine K (**53**) has been detected in the leaves of ferns *Pteridium aquilinum* var. *latiusculum* [58, 65], *Dennstaedtia scaba* [63] and *Hypolepis punctata* [64].

Pteroside K (**54**) is produced as a glucoside by the fern *Pteridium aquilinum* var. *latiusculum* [57].

Pterosine R (**55**) has been found in extracts from *Cibotium barometz* [66]; two other ses-





quiterpenes – (56) and (57) – have been found in *Pteris podophylla* [67] and *Pteridium* subsp. *wightianum* [60], respectively. Histiopterosine B has been isolated from the leaves of fern *Histopterus incisa* [68]. The chemistry and the biological activity of metabolites of indanone type have been described in a review published in 1988 [68].

OTHER TYPES OF SESQUITERPENES

The family *Pluchea* (Asteraceae) is comparatively small and contains 28 genera and approximately 22 species of plants characteristic of South and Central America. Some species have been discovered also in Africa, tropical Asia and Australia [69]. These plants produce sesquiterpenes of eudesmane type among which there are chlorine-containing compounds. For example, two chlorinated derivatives of eudesmane – (59) and (60) – have been found in the plant *Pluchea odorata* widespread in Salvador [70], and two other representatives of this type – (61) and (62) – have been isolated from the leaves of *Pluchea arguta* collected in Pakistan [71]. Out of twelve various sesquiterpenes isolated from the extract of the plant *Laggera crispata* (*Pluchea*) growing in Ethiopia, chlorine is contained in ketoester cuauhtemon (63) [72].

A sesquiterpene of drimane type (64) has been detected in the plant *Makinoa crispata* belonging to the class Bryophyta (liverworts) [73]. Jaeschkenol (65) – a sesquiterpene containing a cyclopropane ring and two chlorine atoms – has been isolated from the plant *Ferula jaeschkeana* (Umbelliferae) growing in the Himalayas (Nepal) [74].

REFERENCES

- J. S. Roberts, in A. A. Newman (Ed.), *Chemistry of Terpenes and Terpenoids*, Acad. Press, London & New York, 1972, p. 88.
- V. M. Dembitsky, G. A. Tolstikov, *Khimiya v interesakh ustoychivogo razvitiya*, 7 (1999) 601.
- V. M. Dembitsky, G. A. Tolstikov, *Ibid.*, 8 (2001) 623.
- G. W. Gribble, *Prog. Chem. Org. Nat. Prod.*, 68 (1996) 1.
- S. M. Kupchan, J. E. Kelsey, M. Maruyama and J. Casady, *Tetrahedron Lett.*, 9 (1968) 3517.
- S. M. Kupchan, J. E. Kelsey, M. Maruyama et al., *J. Org. Chem.*, 34 (1969) 3876.
- J. Harvey-Mason, A. T. Hewson, O. Kennard and R. C. Peterson, *J. Chem. Soc., Chem. Commun.*, (1972) 460.
- J. Lopez de Derma, J. Fayos, S. Garcia-Blanco and M. Martinez-Ripoll, *Acta Crystal.*, 34 B (1978) 2669.
- J. M. Cassaly, D. Abramson, P. Cowall et al., *J. Nat. Prod.*, 42 (1979) 427.
- K. L. Stevens and R. Y. Wong, *Ibid.*, 49 (1986) 833.
- R. I. Evstratova, V. I. Shevchenko, K. S. Rybalko, *Khimiya prirod. soyed.*, (1973) 161.
- K. L. Stevens and R. Y. Wong, *Cryst. Struct. Comm.*, 11 (1982) 949.
- A. G. Gonzalez, J. Bermejo, J. L. Breton and J. Triana, *Tetrahedron Lett.*, 13 (1972) 2017.
- A. G. Gonzalez, J. Bermejo, J. L. Breton et al., *Phytochemistry*, 13 (1974) 1193.
- A. G. Gonzalez, J. Bermejo, J. L. Breton et al., *J. Chem. Soc., Perkin Trans.*, 1 (1976) 1663.
- A. G. Gonzalez, J. Bermejo, J. M. Amaro et al., *Can. J. Chem.*, 56 (1978) 491.
- I. D. Shamyanov, A. Mallabaev, G. P. Sidiyakin, *Khimiya prirod. soyed.*, (1979) 865.
- A. G. Gonzalez, J. Bermejo and G. M. Massanet, *Rev. Latinoamer. Quim.*, 8 (1977) 176.
- S. El-Dahmy, F. Bohlmann et al., *Planta Med.*, 51 (1985) 176.
- J. Jakupovic, Y. Jia, V. P. Pathak et al., *Ibid.*, 52 (1986) 399.
- A. M. Dawidar, M. A. Metwally, M. Abou-Elzahab and M. Abdel-Mogib, *Pharmazie*, 44 (1989) 735.
- D. Youssef and A. W. Frahm, *Planta Med.*, 60 (1994) 267.
- D. Youssef and A. W. Frahm, *Ibid.*, 60 (1994) 573.
- H. S. Al-Easa, J. Mann and A. F. Rizk, *Phytochemistry*, 29 (1990) 1324.
- T. Sarg, S. El-Dahmy, M. El-Domiati and A. Ateya, *Acta Pharm. Hung.*, 58 (1988) 129.
- G. Nowak, M. Holub and M. Budesinsky, *Acta Soc. Bot. Pol.*, 58 (1989) 435.

- 27 G. Nowak, B. Drozdz, M. Holub *et al.*, *Ibid.*, 55 (1986) 227.
 28 G. Nowak, *Phytochemistry*, 31 (1992) 2363.
 29 S. Öksüz, S. Serin and G. Topcu, *Ibid.*, 35 (1994) 435.
 30 A. Rustaiyan, Z. Sharif and A. Tajarodi, *Planta Med.*, 50 (1984) 193.
 31 G. Massiot, A. M. Morfaux, L. LeMen-Oliver *et al.*, *Phytochemistry*, 25 (1986) 258.
 32 S. Öksüz and E. Putun, *Ibid.*, 22 (1983) 2615.
 33 G. Nowak, M. Holub and M. Budesinsky, *Acta Soc. Bot. Pol.*, 57 (1988) 157.
 34 P. Singh and M. Bhala, *Phytochemistry*, 27 (1988) 1203.
 35 M. N. Todorova, I. V. Ognyanov and S. Shatar, *Collect. Czech. Chem. Commun.*, 56 (1991) 1106.
 36 A. A. Mahmoud, A. A. Ahmed and A. A. El-Bassuony, *Fitoterapia*, 70 (1999) 575.
 37 H. Nagaya, T. Nagae and A. Usami, *Nat. Med.*, 48 (1994) 223.
 38 J. A. Marco, R. Sanz, R. Albiach *et al.*, *Phytochemistry*, 32 (1993) 395.
 39 J. Jakupovic, R. Boeker, A. Schuster *et al.*, *Ibid.*, 26 (1987) 1069.
 40 M. I. Yusupov, A. Mallabaev, Sh. Z. Kasymov, G. P. Sidyakin, *Khimiya prirod. soyed.*, 5 (1979) 580.
 41 F. Bohlmann, J. Jakupovic, R. M. King and H. Robinson, *Phytochemistry*, 20 (1981) 1613.
 42 F. Bohlmann, J. Jakupovic, A. Schuster *et al.*, *Ibid.*, 21 (1982) 161.
 43 W. Vichnewski, P. Kulanthaivel, V. L. Goedken and W. Herz, *Ibid.*, 24 (1985) 291.
 44 V. Castro, J. Jakupovic and F. Bohlmann, *Ibid.*, 25 (1986) 1750.
 45 R. Mata, G. Delgado and A. Romo de Vivar, *Ibid.*, 24 (1985) 1515.
 46 H. Wagner, B. Fessler, H. Lotter and V. Wray, *Planta Med.*, 54 (1988) 171.
 47 A. A. Ali, O. M. Abdallah and W. Steglich, *Pharmazia*, 44 (1989) 800.
 48 R. R. Gil, J. A. Pastoriza, J. C. Oberti *et al.*, *Phytochemistry*, 28 (1989) 2841.
 49 A. N. De Gutierrez, E. E. Sigstad, C. A. N. Catalan *et al.*, *Ibid.*, 29 (1990) 1219.
 50 L. V. Reis, M. R. Tavares, F. M. S. B. Palma and M. J. Marcelo-Curto, *Ibid.*, 31 (1992) 1285.
 51 A. A. Ali, N. A. El-Emary, A. A. Khalifa and A. W. Pharm, *Ibid.*, 31 (1992) 2781.
 52 J. Jakupovic, S. Banerjee, V. Castro *et al.*, *Ibid.*, 25 (1986) 1359.
 53 W. Herz and P. Kulanthaivel, *Ibid.*, 22 (1983) 513.
 54 F. Bohlmann, P. K. Gupta, R. M. King and H. Robinson, *Ibid.*, 20 (1981) 743.
 55 F. Bohlmann, P. K. Gupta, R. M. King and H. Robinson, *Ibid.*, 21 (1981) 2593.
 56 K. Yoshihira, M. Fukuoka, M. Kuroyanagi and S. Natori, *Chem. Pharm. Bull. (Japan)*, 19 (1971) 1491.
 57 M. Kuroyanagi, M. Fukuoka, K. Yoshihira and S. Natori, *Ibid.*, 27 (1979) 592.
 58 M. Fukuoka, M. Kuroyanagi, M. Toyama *et al.*, *Ibid.*, 20 (1972) 2282.
 59 J. Kobayashi and K. Koshimizu, *Agric. Biol. Chem.*, 44 (1980) 393.
 60 N. Tanaka, T. Satake, A. Takahashi *et al.*, *Chem. Pharm. Bull. (Japan)*, 30 (1982) 3640.
 61 T. Kuraishi, T. Murakami, T. Taniguchi *et al.*, *Ibid.*, 33 (1985) 2305.
 62 T. Murakami, H. Maehashi, N. Tanaka and T. Satake, *J. Pharmacol. Soc. Japan*, 105 (1985) 640.
 63 T. Murakami, K. Owashi, N. Tanaka *et al.*, *Chem. Pharm. Bull. (Japan)*, 23 (1975) 1630.
 64 Y. Hayashi, M. Nishizawa, S. Harita and T. Sakan, *Chem. Lett.*, (1972) 375.
 65 M. Fukuoka, M. Kuroyanagi, K. Yoshihira and S. Natori, *Chem. Pharm. Bull. (Japan)*, 26 (1978) 2365.
 66 T. Murakami, T. Satake, K. Ninomiya *et al.*, *Phytochemistry*, 19 (1980) 1743.
 67 N. Tanaka, T. Murakami, Y. Saiki *et al.*, *Chem. Pharm. Bull. (Japan)*, 29 (1981) 3455.
 68 T. Murakami and N. Tanaka, *Prog. Chem. Org. Nat. Prod.*, 54 (1988) 1.
 69 A. A. Anderberg, in K. Bremer (Ed.), *Asteraceae: Cladistic and Classification*, Timber Press, Portland, Oregon, 1995, p. 301.
 70 F. J. Arriaga-Giner, J. Borges Del Castillo, M. T. Mansilla-Ferrero *et al.*, *Phytochemistry*, 22 (1983) 1767.
 71 V. U. Ahmad, T. A. Farooqui, K. Fizza *et al.*, *J. Nat. Prod.*, 55 (1989) 730.
 72 A. A. Ahmed, H. R. El-Seedi, A. A. Mahmoud *et al.*, *Phytochemistry*, 49 (1998) 2421.
 73 T. Hashimoto, M. Tori and Y. Asakawa, *Ibid.*, 28 (1989) 3377.
 74 S. N. Gaeg, S. K. Agaewal, K. Fidels *et al.*, *J. Nat. Prod.*, 56 (1993) 539.